







Erratum: Prevalence and distribution of selected cervical human papillomavirus types in HIV infected and HIV uninfected women in South Africa, 1989–2021: A narrative review

**Authors:**

Rixongile R. Rikhotso¹ 
Emma M. Mitchell² 
Daniel T. Wilson³ 
Aubrey Doede⁴ 
Nontokozo D. Matume⁵ 
Pascal O. Bessong^{5,6} 

Affiliations:

¹Department of Biochemistry and Microbiology, Faculty of Science, Engineering and Agriculture, University of Venda, Thohoyandou, South Africa

²Department of Family, Community and Mental Health Systems, School of Nursing, University of Virginia, Charlottesville, United States of America

³Claude Moore Health Sciences Library, School of Nursing, University of Virginia, Charlottesville, United States of America

⁴Department of Family Medicine and Public Health, University of California San Diego, California, United States of America

⁵HIV/AIDS & Global Health Research Programme, University of Venda, Thohoyandou, South Africa

⁶Center for Global Health Equity, School of Medicine, University of Virginia, Charlottesville, Virginia, United States of America

Corresponding author:

Rixongile Rikhotso,
rikhotsorrrr@gmail.com

Read online:

Scan this QR code with your smart phone or mobile device to read online.

In the published article, Rikhotso RR, Mitchell EM, Wilson DT, Doede A, Matume ND, Bessong PO. Prevalence and distribution of selected cervical human papillomavirus types in HIV infected and HIV uninfected women in South Africa, 1989–2021: A narrative review. *S Afr J Infect Dis.* 2022;37(1), a363. <https://doi.org/10.4102/sajid.v37i1.363>, an omission error occurred in Online Appendix 1 (supplementary data). Table 2, Table 3 and Table 4, were initially omitted from Online Appendix 1 on the first publication. The updated and correct Online Appendix 1 showing all four tables replaced the incorrect version with its republication.

The publisher apologises for this error. The correction does not change the study's findings of significance, the overall interpretation of the study's results, or the scientific conclusions of the article in any way.

Dates: Published: 30 Jan. 2024

How to cite this correction: Rikhotso RR, Mitchell EM, Wilson DT, Doede A, Matume ND, Bessong PO. Erratum: Prevalence and distribution of selected cervical human papillomavirus types in HIV infected and HIV uninfected women in South Africa, 1989–2021: A narrative review. *S Afr J Infect Dis.* 2024;39(1), a575. <https://doi.org/10.4102/sajid.v39i1.575>







Copyright: © 2024. The Authors. Licensee: AOSIS. This work is licensed under the Creative Commons Attribution License.

Note: DOI of original article published: <https://doi.org/10.4102/sajid.v37i1.363>

Prevalence and distribution of selected cervical human papillomavirus types in HIV infected and HIV uninfected women in South Africa, 1989–2021: A narrative review



Authors:

Rixongile R. Rikhotso¹ 
 Emma M. Mitchell² 
 Daniel T. Wilson³ 
 Aubrey Doede⁴ 
 Nontokozo D. Matume⁵ 
 Pascal O. Bessong^{5,6} 

Affiliations:

¹Department of Biochemistry and Microbiology, Faculty of Science, Engineering and Agriculture, University of Venda, Thohoyandou, South Africa

²Department of Family, Community and Mental Health Systems, School of Nursing, University of Virginia, Charlottesville, United States of America

³Claude Moore Health Sciences Library, School of Nursing, University of Virginia, Charlottesville, United States of America

⁴Department of Family Medicine and Public Health, University of California San Diego, California, United States of America

⁵HIV/AIDS & Global Health Research Programme, University of Venda, Thohoyandou, South Africa

⁶Center for Global Health Equity, School of Medicine, University of Virginia, Charlottesville, Virginia, United States of America

Corresponding author:

Rixongile Rikhotso,
rikhotsorrr@gmail.com

Read online:



Scan this QR code with your smart phone or mobile device to read online.

Background: Human papillomavirus infection, a causative agent of cervical cancer, is of great concern, more so in populations with high HIV prevalence, such as South Africa.

Aim: This review aimed to examine the prevalence and distribution of selected cervical human papillomavirus (HPV) types in HIV infected and HIV uninfected women in South Africa.

Methods: PubMed and Web of Science databases were searched using key words. For data integrity, data was assessed by two authors independently. The study inclusion criteria comprised records on cervical HPV, HPV genotyping and HPV type distribution among South African women. Statistical analysis was performed using Social Science Statistics.

Results: Sixty-nine articles met the inclusion criteria for analysis. Data on cervical HPV prevalence and type distribution was available only for five of the nine provinces of South Africa. Only 4/69 studies used sequencing as an approach to identify HPV types. In a general population, HPV type 16 was the most frequent (8.80%), followed by types 35 (4.86%), 18 (4.14%), 58 and 52 with the frequency of 3.65% and 3.62%, respectively. Furthermore, the least frequent type was HPV 70 (0.74%). Both HIV infected and HIV uninfected populations had a higher prevalence of high-risk human papillomavirus (hrHPV) types 16, 18 and 35 than other HPV types; while HPV types 6, 11 and 70 were the least frequent types from these populations. Lastly, HPV 16 was the most predominant type among women with normal (2.03%) and abnormal cervical cytology (6.60%).

Conclusion: Expanding on HPV genotyping will improve the knowledge in patterns of HPV type distribution in South Africa that will further help in decision making to improve current diagnostics, and future vaccine development and assessment.

Keywords: HPV types; HPV prevalence; HPV; HIV; Women; South Africa.

Introduction

Human papillomavirus (HPV) and human immunodeficiency virus (HIV) are leading sexually transmitted infections (STIs)¹ that cause significant burden of disease in low- and middle-income countries.^{2,3} Cervical cancer (CC) is underlined as an HIV-associated malignancy and is rated as the second most frequent cancer among South African women of all age groups⁴ after breast cancer. About 70% of CC cases are caused by persistent infection of high risk human papillomavirus (hrHPV) types 16 and 18.⁵ In 2020, there were 10 702 new cases of CC among South African women reported, irrespective of HIV status; of these 5870 women died because of the disease.⁶

Globally, South Africa has the highest HIV infection rates with an estimate of 8.23 million people living with HIV in 2021⁷ with women being the most affected group. However, the introduction and free access to anti-retroviral therapy (ART) has increased the life span of individuals living with HIV and AIDS,⁸ leading to an increasing number of people ageing with HIV infection. Women living with HIV subsequently experience life expectancy approaching that of persons

Dates: Received: 27 Oct. 2021 | Accepted: 10 Mar. 2022 | Published: 08 June 2022

How to cite this article: Rikhotso RR, Mitchell EM, Wilson DT, Doede A, Matume ND, Bessong PO. Prevalence and distribution of selected cervical human papillomavirus types in HIV infected and HIV uninfected women in South Africa, 1989–2021: A narrative review. *S Afr J Infect Dis.* 2022;37(1), a363. <https://doi.org/10.4102/sajid.v37i1.363>

Copyright: © 2022. The Authors. Licensee: AOSIS. This work is licensed under the Creative Commons Attribution License.

Note: Additional supporting information may be found in the online version of this article as Online Appendix 1.

living without HIV, although, they still have an increased incidence of developing CC at a later stage⁹ probably due to their compromised immune system.

According to the International HPV Reference Center, to date there are 228 HPV genotypes that are known,¹⁰ and about 40 of these genotypes have been reported to infect the anogenital region.¹¹ Based upon the potential interconnection between various HPV genotypes and CC, HPV was thereafter stratified into two main groups: low (lr) and high risk (hr) genotypes primarily known to be non-cancerous¹² and cancerous,¹³ respectively.

There is currently no cure for HPV infection. However, prevention is possible through vaccination. Three HPV vaccines have been developed thus far, which offers protection against some of the hr types. The global uptake of these vaccines varies geographically.⁹ Cervarix is a bivalent vaccine that covers HPV types 16 and 18; Gardasil is a quadrivalent vaccine offering protection against HPV genotype 6, 11, 16 and 18. Lastly, Gardasil-9 is a nonavalent vaccine and comprises the four HPV types covered by the quadrivalent vaccine, with the addition of types 31, 33, 45, 52, and 58. However, in South Africa, only Cervarix and Gardasil are currently available.

The Department of Health in collaboration with the Department of Basic Education launched a South African HPV vaccination campaign in 2014. The campaign specifically targets 9-year olds and above public-school children in grade 4 and involved delivering two doses of vaccine 6 months apart.^{14,15} In 2014, it was reported that more than 350 000 school girls received the vaccine which covered more than 16 000 of public schools. Within the same year of the programme, 86.6% coverage was attained.¹⁴ These campaigns and programmes play a role in the reduction of HPV infections and effectively control and prevent CC development in the later stage. However, issues of cost and unavailability of vaccines limit the progress of these programmes as well as the limitation of the available HPV vaccines to cover all the prevalent HPV or hrHPV types.

Immunocompromised individuals, especially HIV-positive women are at high risk of HPV infections¹⁶ and associated pathologies. Studies have shown a high prevalence of hrHPV types in HIV positive women than negative women^{17,18} and other studies indicate multiple hrHPV infections in HIV positive women.^{19,20} However, other studies indicate a high prevalence of HPV infections in women regardless of HIV status.^{19,21} Therefore, there are uncertainties on the prevalence and distribution of HPV types regarding HIV status, and more studies are needed to understand which hrHPV types are associated with HIV and how they contribute to the development of CC in this population, especially in South Africa where there is a high prevalence of HIV infection.

For these reasons, this review reports on the literature specific to South Africa, highlighting the following: (1) The distribution of cervical HPV studies across provinces in

South Africa, (2) The diagnostic methods used to identify HPV types in South Africa, (3) The prevalence and distribution of hrHPV types among women in general, and among those with a normal and abnormal cervical cytology, (4) The prevalence and distribution of hr/lrHPV types by HIV status.

Review methodology

Eligibility criteria

Research articles focusing on cervical HPV infection in South Africa were sought. Records on cervical HPV, HPV genotyping and type distribution published among South African women formed part of the study inclusion criteria. Five major exclusion criteria were designated: (1) not geographically related (South Africa must have been at least one of the countries included); (2) review articles, clinical trials and preface; (3) poster presentations, letters to the editor and correspondence; (4) not HPV or cervical (HPV or CC) related and (5) HPV in men. Full text articles published in English were included. Of note, comparative studies conducted between South Africa and other countries were included in the final analysis. However, in such studies only South African data relevant to our review was extracted.

Information sources

Published reports on the prevalence and cervical HPV types among South African women living with and without HIV were carefully searched and compiled. In addition, articles that did not differentiate HPV prevalence and types by HIV status were also included to answer objectives which were independent of HIV status. All the reports were identified through PubMed and Web of Science databases last consulted on 01 July 2021.

Search strategy

All the records were sourced using an advanced search option for both databases. In addition, all fields were also searched. The following search strategies were generated per database:

PubMed: (((HPV) OR ((Human Papillomavirus))) OR ((Human Papilloma virus))) OR (Papillomaviridae) AND ((South Africa*))

("HPV"[All Fields] OR ("alphapapillomavirus"[MeSH Terms] OR "alphapapillomavirus"[All Fields] OR ("human"[All Fields] AND "papillomavirus"[All Fields]) OR "human papillomavirus"[All Fields]) OR ("papillomaviridae"[MeSH Terms] OR "papillomaviridae"[All Fields] OR ("human"[All Fields] AND "papilloma"[All Fields] AND "virus"[All Fields]) OR "human papilloma virus"[All Fields]) OR ("papillomaviridae"[MeSH Terms] OR "papillomaviridae"[All Fields])) AND ("South"[All Fields] AND "africa*" [All Fields])

Translations

Human papillomavirus

"alphapapillomavirus"[MeSH Terms] OR "alphapapillomavirus"[All Fields] OR ("human"[All Fields] AND

“papillomavirus”[All Fields] OR “human papillomavirus” [All Fields]

Human papilloma virus

“papillomaviridae”[MeSHTerms]OR“papillomaviridae”[All Fields] OR (“human”[All Fields] AND “papilloma”[All Fields] AND “virus”[All Fields]) OR “human papilloma virus”[All Fields]

Papillomaviridae

“papillomaviridae”[MeSHTerms]OR“papillomaviridae”[All Fields]

Web of science

TS=(HPV) OR TS=(Human Papillomavirus) OR TS=(Human Papilloma virus) OR TS=(Papillomaviridae) AND TS=(South Africa)

Selection process and study risk of bias assessment

The full-text of the relevant articles were read and analysed for the objectives of the current study. Variables required to answer the study objectives were extracted and tabulated into their respective columns. Two authors independently reviewed the search and selection procedure for consistency. Of note, the authors agreed that only articles that strictly speak of the study inclusion criteria be the ones to be considered for further analysis. Variables required that were not reported in certain articles were marked and the authors agreed that such information should be labelled as ‘data was not reported’.

Data collection process


All records searched from both databases were then exported to Mendeley Desktop, a referencing tool and further processed to remove duplicates and screened for eligibility. The title and abstract were thoroughly screened in Mendeley Desktop and later full articles were downloaded and screened one at a time. This was done by two authors independently using the same tool.

Data items

Research articles focusing on cervical HPV infection in South Africa published between 1989 and 2021 were retrieved. Data extracted included: (1) relevant information on sample type (cervical cytology/cervicovaginal, biopsies) and from women; (2) sampling device and genotyping technique used; (3) study design, collection time period and age range; (4) study population and sample size; (5) study setting; and (6) data on cervical status. Following the PRISMA protocol,²² (Figure 1) a total of 69 published articles were included for final analysis after screening.

Synthesis method

The full-text of the relevant articles were read and analysed for the objectives of the current study which were grouped



Source: Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. PLoS Med. 2009;6(7):e1000097. <https://doi.org/10.1093/ptj/89.9.873>

HPV, human papillomavirus.

FIGURE 1: Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) workflow for the selection of studies to examine the human papillomavirus types among South African women living with and without human immunodeficiency virus (1989–2021).

into four objectives: (1) The distribution of cervical HPV studies across provinces in South Africa, (2) The diagnostic methods used to identify HPV types in South Africa, (3) The prevalence and distribution of hrHPV types among women in general population and among those with a normal and abnormal cervical cytology, (4) The prevalence and distribution of hr/lrHPV types by HIV status.

To achieve objective 1 and 2, variables required to answer these objectives were extracted from individual records and tabulated into their respective columns using a word file. The variables were (1) author (year), and study title, (2) sample type, sampling device and genotyping technique, and (3) province in South Africa. Of note, these objectives were analysed regardless of the HIV status of study participants recruited, considering all the 69 studies that were included in the quantitative analysis. Records missing certain required information were identified and marked as not reported (Supplementary Table 2).

To achieve objective 3, data was first pooled from 46 of 69 studies to estimate the prevalence and distribution of hrHPV types among women in the general population that is regardless of the HIV status in the country (Supplementary Table 3). To do so, the 69 records were further scrutinised and

all the records that (1) reported the distribution of HPV types based on the vaccine coverages, (2) reported HPV types collectively as lr/hr HPV types and/or single/multiple infections instead of reporting on the individual types across the study participants, (3) studies that updated data from the previously published data, (4) studies whose results were showed by graphs making it difficult to interpret the proportions based on the scales used, and (5) comparative studies which did not separate the distribution of the HPV DNA by country, such records were excluded. Of note, to estimate the overall number of study participants, studies that used the same samples such participants were counted once, this was done to avoid over-representation of the population size. Furthermore, a sub-analysis was conducted from 22/46 studies that clearly documented the prevalence of hr-HPV types among South African women according to cervical cytology status.

To achieve objective 4, which was to determine the prevalence and distribution of hr/lrHPV types by HIV status, an intra-search strategy using Mendeley was conducted again within the 46 studies. The following steps were taken. The upper right section of Mendeley page was clicked to help segregate the records by HIV status. Key words such as Human Immunodeficiency Virus, HIV, HIV-infected/positive and HIV-uninfected/negative/HIV non-infected and HIV non-positive were entered into the search box interchangeably to search for HPV records in the library that speaks of HIV status. The resulting 26 records that clearly specified the distribution of HPV types according to HIV status in the country were then grouped into a sub-folder for analysis. Of note, part of the excluded records included all studies that had unreported South African province, unspecified HIV status and those that did not segregate the distribution of HPV types by HIV status.

Analysis for human papillomavirus types prevalence and distribution in South Africa

Eligible literature was screened by provinces in South Africa that reported for HPV types, types of specimen, laboratory technique used to identify HPV types and HIV status.

Outcomes were tabulated according to author and year of publication, province of study, study population (women living with and without HIV), and study design. Extracted data was examined, across the country, for the prevalence of each of the 14 hrHPV types (16, 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, 68 and 70)^{23,24} regardless of the HIV status first and then classified according to HIV status with two low risk human papillomavirus (lrHPV) type 6 and 11 included.

Statistical analysis

An online Social Science Statistics tool using option *p*-value calculators was used to derive the *p*-values from the chi square. This was employed in order to determine the association between HIV status and the distribution of

HPV infection. A positive significant test was considered statistically significant at $p < 0.05$.

Review findings

Characteristics of studies analysed

According to our inclusion criteria, 69 studies were selected for analysis, and represented only five (Western Cape, Eastern Cape, Gauteng, KwaZulu-Natal and Limpopo) of the nine provinces of South Africa. Overall, a total of 38 049 women participated in these studies, of whom, 9955 and 20 747 were living with and without HIV, respectively. Western Cape province had the highest number of study participants ($n = 27\ 779$) followed by Gauteng province ($n = 5817$), Limpopo province has been observed to have the least number ($n = 100$) of study participants. Of note, to avoid over representation of the study participants, all studies that used the same sample size were counted once. Supplementary Table 1, show the characteristics of all the 69 articles included in the quantitative analysis.

Fifty two of the 69 studies were conducted uniquely in Western Cape ($n = 31$), Gauteng ($n = 12$), Eastern Cape ($n = 5$), KwaZulu-Natal ($n = 3$), and Limpopo ($n = 1$). In two studies, participants were recruited from more than one province: one study comprised participants from KwaZulu-Natal and Gauteng, and the other had participants from Western Cape and Gauteng provinces. In two studies, the province from which participants were recruited was not indicated. Furthermore, 13 comparative studies were conducted between South Africa and other countries: three studies were reported for Britain and South Africa (KwaZulu-Natal), seven studies reported for Burkina Faso and South Africa (Gauteng), and one study was conducted in three countries – Ghana, Nigeria, and South Africa. Lastly, one study each was conducted in Limpopo and Tanzania while the other one in KwaZulu-Natal and Nairobi. These studies are represented in Supplementary Table 2.

The 69 studies eligible for analysis comprised three study populations: HIV non-infected ($n = 6$), HIV-positive ($n = 17$), and both HIV positive and non-infected populations ($n = 31$). Fifteen studies had unspecified HIV status. Overall, there were 48 studies that included HIV-positive populations, including those that comprised both HIV-positive and HIV uninfected individuals.

Sample type, collection mode, and human papillomavirus genotyping approaches

Of the 69 studies analysed, we grouped specimens into cervical/cervicovaginal lavages, biopsies, self-collected and both self-collected/cervical specimens. The most commonly used specimens included cervical/cervicovaginal lavages, reported in 40 studies, and cervical biopsies, reported in 17 studies. In addition, in six studies specimens were both cervical and self-collected, while in six studies self-collected specimens were used. Of the devices and tools used to collect these specimens, brushes were the commonly used tool reported in 32 of the 69 studies. The use of a

spatula was among the least used collection device, Supplementary Table 2.

There is a fair degree of heterogeneity on the approaches used to genotype HPV. Linear Array (LA) was the most commonly used method reported in 30 studies. This was followed by INNOLiPA ($n = 6$) and hybrid capture 2 (HC2) $n = 4$. Enzyme immuno assay (EIA), non-isotopic *in situ* hybridisation (NISH), and Xpert HPV were used in three studies each, while southern blot hybridisation was applied in two studies. HPV direct flow chip and TaqMan based quantification polymerase chain reaction (PCR) was applied once, respectively. Of the studies that used more than one technique, HC2/LA was the most commonly used ($n = 4$), followed by HC2/INNO Lipa ($n = 2$). The least frequently used technique included LA/Xpert HPV, care HPV/INNO Lipa, reverse line blot/PreTeck HPV proofer, HC1/HC2, HC2/hpVIR, and cobas test/LA/HC2 which were used in one study each. In one comparative study, Illumina detected 16 HPV types, including type 30, 74, 86 and 90, not included in LA; and Sanger sequencing was seen to resolve HPV DNA (Deoxyribonucleic acid) untypable by Restriction fragment length polymorphism (RFLP) in another study. Lastly, one study used southern blot hybridisation and sequencing, while the other study compared the genotypic outcomes obtained through massive parallel sequencing to HPVIR assay LA HPV genotyping test (Supplementary Table 2).

High risk human papillomavirus type diversity and human immunodeficiency virus status

Firstly, the proportion of 14 hrHPV types was examined, irrespective of the HIV status of the women. Overall, 46 studies were thereafter examined, for the 14 hrHPV types. Of a total of 30 677 participants, HPV type 16 was the most frequent (8.80%), followed by types 35 (4.86%), 18 (4.14%), 58 and 52 with the frequency of 3.65% and 3.62% individually. The least frequent was type 70 (0.74%), (Supplementary Table 3).

Furthermore, a sub-analysis of cytology data demonstrated that a total of 13 044 and 15 782 women had a normal and abnormal pap smear results, respectively. Women with abnormal pap smear had higher prevalences of HPV types when compared to their counterparts. Among women who had an abnormal pap smear, HPV types 16 (6.60%), 35 (2.43%), 18 (2.24%), 58 (1.70%), 33 (1.62%), 45 (1.57% each) and 52 (1.52%) were the seven most prevalent types in descending order of frequency. On the contrary, HPV types 16 (2.03%), 35 (1.85%), 58 (1.58%), 45 (1.55%), 18 (1.40%), 52 (1.30%) and 68 (1.19%) were the most predominant types among women who had a normal pap smear test.

From the 26 studies that clearly documented the proportion of HPV type distribution according to HIV status, these studies were examined to estimate the frequency of 14 hrHPV and 2 lrHPV types and to determine the statistical association between HPV type distribution and HIV status, also

considering the types included in Gardasil-9 vaccine. The findings revealed that, similar HPV types were detected in both HIV-positive and HIV-uninfected women, with HPV 16, 18 and 35 being the most common detected types in both HIV-positive and HIV-uninfected populations in the country; however, the proportions varied (Supplementary Table 4). For these types, the proportions observed were as follows: 0.135% and 0.047% (HPV 16); 0.065% and 0.028% (HPV 35); 0.068% and 0.024% for type 18 detected in women living with and without HIV, respectively. On the contrary, HPV 6, 11 and 70 were the least common types with the proportions ranging from 0.001% to 0.015%. Furthermore, statistical analysis depicted that types 6, 11, 16, 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, 68 and 70 were significantly more common among women living with HIV than those who were living without HIV, while type 59 was the only type observed to be significantly more common among women living without HIV. All these HPV types were statistically significant at $p < 0.00001$ (Supplementary Table 4).

Implications, recommendations and conclusion

This review provides insights on the scope of HPV research on viral diversity and HIV status in South African women between 1989 and 2021. Primarily, it reveals that HPV research is limited to five (Gauteng, KwaZulu-Natal, Limpopo, Western Cape and Eastern Cape) of the nine provinces of the country, and even in this regard, there are relatively few studies from each of the five provinces. Nevertheless, Western Cape province was observed to have the highest number of studies among the five provinces. Factors such as poor access of primary healthcare systems, lack of viral HPV knowledge, lack of pap testing, misconceptions and conspiracy theories associated with the HPV infection at large could have had an impact on the participation of the participants in the studies. Factors such as financial constraints involved to carry out the HPV diversity research studies, lack of HPV research expertise, difficulties in attaining permission (s) and approval (s) from relevant organisations might have been some of the reasons why other provinces lack data on HPV research.

Studies have shown that the distribution of HPV infection differs by province, technique used to detect it, HIV status of participants, specimen collection method and the age group of participants.^{25,26,27,28} The study findings revealed that there are various methods used to identify HPV types in the country, with Roche LA being the most commonly used technique while sequencing is rarely used. Thus, more consistent methods for sample collection and genotyping procedures are necessary to systematically assess the prevalence of potentially carcinogenic types of HPV.

Despite the higher cost involved in next generation sequencing (NGS),²⁹ the genotyping tool has proved to be most specific³⁰ and sensitive^{31,32} compared to LA genotyping test kit that only detects the restricted number of 37 HPV types incorporated in the kit and has the possibility of

missing mixed and minority infections. Flores-Miramontes et al. in Mexico genotyped HPV using both LA and NGS, detected HPV types 32, 44, 74, 102, and 114 not incorporated in the LA.³² In South Africa, specifically Cape-Town, Meiring et al. compared LA and NGS and found that NGS was able to detect four additional HPV types (30, 74, 86 and 90) not covered by the 37 restricted types of the LA.³³ This also implies that the differences on how the samples are genotyped have a great impact on the genotyping outcomes. Thus, the authors agree that more sensitive technologies such as NGS are significant at least at a research level to be used as a strategy to unravel the full spectrum of HPV types in the South African population. Of interest, recent findings documented by Ardhaoui and colleagues, depicted that NGS was able to detect genotypes of clinical importance that include HPV type 31, 39 and 52 that the reverse line hybridisation genotyping technique failed to detect regardless of the presence of their probes embedded on the membrane.³⁴ This raises a significant aspect of presumably under reporting of other HPV types of clinical importance in certain cohorts. Of note, kits validation for accuracy purposes and in the management of the disease can be encouraged.

Despite the inconsistencies used in data reporting strategies across the studies, the overall HPV type distribution according to HIV status and in a general population depict that HPV 16, 35 and 18 were the most frequent types in these populations; thus, encourages the need for routine hospital or clinic visits for regular check-ups among South African women, especially those who never received any vaccine. With evidence pointing to a significant increase of HPV more in women living with HIV^{16,35,36,37,38} and in the general population. The findings of the literature review conducted in 2021 by Kombe and colleagues highlighted that, the prevalence of HPV type distribution do vary across the world although that does not change the associated diseases caused by these viral types. Furthermore, there are certain HPV types that have been found to be present around the world.³⁹ Several review studies conducted across continents including Africa, revealed that hrHPV types such as 16, 18, 35 and 58 among others are the most frequent types³⁹ which validates our study findings.

Human papillomavirus/human immunodeficiency virus co-infection among women particularly in developing countries remains one of the principal public health challenges requiring closer intervention. Worldwide, in a general population HPV types 16, 18 and 45 are primarily linked with CC²³; however, the prevalence of HPV types varies worldwide. In Asia, HPV 52 and 45 are the most prevalent HPV types; in the USA and Europe, HPV types 45 and 33, respectively, are the most common types, and types 35 and 45 are more predominant in Africa,^{40,41,42} all these types were common after HPV 16 and 18 in participants with invasive CC. Both HPV types 35 and 45 are considered hrHPV²³ for subsequent CC. Although a number of studies show that the bivalent vaccine does offer protection against these types, cross protection vaccines are less reliable compared to type specific and is of a limited duration.⁴³

Neither of these are included in the currently available HPV vaccines in South Africa. To enrich our understanding of HPV types, including detection of minority viral populations in South Africa, the application of sanger sequencing or next-generation sequencing is strongly recommended,⁴⁴ at least for research purposes. This is important to identify putative hrHPV types which may not be covered by the current diagnostic approaches and vaccines.

The state of cervical cytology in the studied population plays a significant role in the distribution of HPV types, be it single and/or multiple infections and the overall prevalence. Our findings revealed that HPV types 16, 35, 58, 45, 18, 52 and 68 and types 16, 35, 18, 58, 33, 45 and 52 were the most prevalent types in descending order among women with normal and abnormal pap smear, respectively. This was in accordance with a study conducted by Wolday and colleagues on the distribution of HPV genotype among Ethiopian women with normal and abnormal cervical cytology. Their findings revealed that HPV types 16, 35, 45, and 18 were among the most prevalent HPV types among women with normal cytology while most women with abnormal cervical cytology had HPV types 16, 45, 31 and 35.⁴⁵ The findings of another study conducted in Turkey by Muderris and colleagues agree with that of our study findings.⁴⁶ In both studies HPV type 16 was the most detected type in both compartments. A meta-analysis data by Obeid et al.⁴⁷ demonstrated that HPV 16 and 58 were the most common types among women with abnormal cervical cytology in the Middle East and North Africa. Of note, we noticed that a normal cervical cytology test does not necessarily mean the absence of hrHPV of clinical importance. Thus, the presence of hrHPV types among women with normal cytology raises a serious challenge among policymakers in South Africa. The study findings encourage that more closer routine check-ups and HPV screening should be prioritised among South African women.

Cervical cancer is responsible for significant mortality among women.³² Cervical cancer mortality among women living with HIV infection is twice greater than those living without HIV.^{48,49} Nonetheless, the availability of ART increases the life span of women living with HIV and raises a necessity for the prevention of CC in developing countries.⁵⁰ To fully understand the distribution of HPV types in a specific geographic location, genotyping samples from both individuals living with HIV and without HIV is highly recommended. In the absence of effective screening, CC might kill millions of women.⁵¹ Human papillomavirus genotyping can be both a secondary prevention technique employed to help in the screening and early detection of CC, as well as a tertiary prevention technique that also helps in the management of pre-cancerous cervical lesions.⁵² In addition, HPV genotyping technique can be used to further investigate women who test positive during cytology.⁵³ Possible recurrence of HPV infections can also be monitored using this method during patient follow-ups after their treatment of cervical intraepithelial neoplasia (CIN).^{54,55} Thus, understanding the facet of HPV types would aid in the better strategies of prevention and management of CC.⁵² Despite the

late HPV vaccination rollout in South Africa, Western Cape, Gauteng and KwaZulu-Natal provinces do have data depicting the vaccination coverages of first doses among the primary schools ranging between 64.0% and 99.7%.^{56,57,58} However, to the best of our knowledge, checking when the vaccination rollout programme commenced in the country and the target age groups, our study participants analysed are highly unlikely to have been vaccinated or benefited from these programmes. Therefore, perhaps some of the circulating types of clinical significance might have been prevented or lowered provided some of the study participants benefited from the vaccine.

This narrative review should be considered with some limitations. It was not possible to look at the differences in the distribution of HPV types according to socio-economic and vaccine status of the participants due to the lack of data in the analysed studies. It has been demonstrated that these two factors impact the carriage and type of HPV.^{59,60} Secondly, it would have been of interest to review the distribution of hrHPV in women on antiretroviral therapy. Recent data suggests that ART may lead to clearance of HPV infection.^{61,62} Lastly with South Africa having high levels of disparities in terms of access to health and health outcomes from rural and urban settings, it would have been great to determine the HPV type distribution and prevalence in those two settings. However, out of 69^{63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100,101,102,103,104,105,106,107,108,109,110,111,112,113,114,115,116,117,118,119,120,121,122,123,124,125} studies that met our inclusion criteria only six (9%) were conducted specifically in rural areas. Thus, a comparison would be biased towards where more studies were done.

In conclusion, this review describes the prevalence and distribution of cervical HPV types as reported in the literature which only represents participants recruited from five of nine provinces in South Africa, with Roche LA being the commonly used genotyping technique. This raises a potentially serious concern due to the lack of research studies and data in the country. Furthermore, we observed that hrHPV type 16 was the most predominant type from all the populations analysed. Therefore, further research to expand the scope of HPV types is proposed, including the use of sensitive approaches such as NGS. This will augment the understanding of the patterns of HPV geo-diversity in South Africa, that will further help in decision making with regard to study vaccines and screening strategies that are geographically relevant.

Acknowledgements

The authors would like to thank the South African National Research Foundation and Fogarty International Center of the NIH, USA for funding this study.

Competing interests

EMM has received research funding from Merck, Inc. in the last 5 years. RRR, AD, DW, NDM, and POB have no conflicts of interest to disclose.

Authors' contributions

P.O.B., E.M.M., and R.R.R. were responsible for the conceptualisation and design of the study; R.R.R., E.M.M. and D.W. did the data collection and assembly of data; P.O.B., R.R.R., N.D.M., E.M.M. were responsible for the data analysis and interpretation; All authors were responsible for the manuscript writing; A.D., R.R.R., N.D.M., E.M.M. did the tables and data visualisation; All authors were responsible for the final approval of the manuscript: all authors agreed to be accountable for all aspects of the work.

Ethical considerations

This article followed all ethical standards for research without direct contact with human or animal subjects.

Funding information

The study was supported by the South African National Research Foundation grants to POB (GUN113465, GUN109312, GUN86037). Additional support was provided by the Fogarty International Center of the NIH, USA under Award Number D43 TW006578 to RRR and NDM.

Data availability

Data sharing is not applicable to this review article as no new data were created or analysed in this study.

Disclaimer

The content is solely the responsibility of the authors and does not necessarily represent the official views of the South African National Research Foundation, Fogarty International Center or the NIH.

References

- Lissouba P, Van de Perre P. Association of genital human papillomavirus infection with HIV acquisition: A systematic review and meta-analysis. *Sex Transm Infect.* 2013;89(5):350–356. <https://doi.org/10.1136/sextrans-2011-050346>
- Moodley JR, Hoffman M, Carrara H, et al. HIV and pre-neoplastic and neoplastic lesions of the cervix in South Africa: A case-control study. *BMC Cancer.* 2006;6(1):135. <https://doi.org/10.1186/1471-2407-6-135>
- De Sanjosé S, Serrano B, Castellsagué X, et al. Human papillomavirus (HPV) and related cancers in the Global Alliance for Vaccines and Immunization (GAVI) countries. A WHO/ICO HPV Information Centre Report. *Vaccine.* 2012;30(4):D1–D83. [https://doi.org/10.1016/s0264-410x\(12\)01435-1](https://doi.org/10.1016/s0264-410x(12)01435-1)
- Bruni LB, Barrionuevo-Rosas L, Albero G, et al. Human papillomavirus and related diseases in the world. Barcelona: ICO information Centre on HPV and Cancer (HPV information Centre). Summary Report; 2015.
- Reid R, Stanhope CR. Genital warts and cervical cancer. I. Evidence of an association between subclinical papillomavirus infection and cervical malignancy. *Cancer.* 1982;50(2):377–387. [https://doi.org/10.1002/1097-0142\(19820715\)50:2%3C377::aid-cnrcr2820500236%3E3.0.co;2-a](https://doi.org/10.1002/1097-0142(19820715)50:2%3C377::aid-cnrcr2820500236%3E3.0.co;2-a)
- Bruni L, Albero G, Serrano B, et al. Human papillomavirus and related diseases in South Africa. Barcelona: ICO/IARC Information Centre on HPV and Cancer (HPV Information Centre). Summary Report; 2021.
- Statistics South Africa. Mid-year population estimates 2021 [homepage on the Internet]. [cited 2022 Feb 02] Available from: www.statssa.gov.za
- Hogg R, Lima V, Sterne JAC, et al. Life expectancy of individuals on combination antiretroviral therapy in high-income countries: A collaborative analysis of 14 cohort studies. *Lancet.* 2008;372:293–299. [https://doi.org/10.1016/s0140-6736\(08\)61113-7](https://doi.org/10.1016/s0140-6736(08)61113-7)
- Loke AY, Kwan ML. The uptake of human papillomavirus vaccination and its associated factors among adolescents: A systematic review. *J Prim Care Community Health.* 2017;8(4):349–362. <https://doi.org/10.1177/2150131917742299>

10. Karolinska Institutet. International Human Papillomavirus Reference Center. [cited 2021 July 07]. Available from: <http://www.hpvcenter.se>
11. De Villiers EM, Fauquet C. Classification of papillomaviruses. *Virology*. 2004;324(1):17–27. <https://doi.org/10.1016/j.virol.2004.03.033>
12. Bhatia N, Lynde C. Understanding genital warts: Epidemiology, pathogenesis, and burden of disease of human papillomavirus. *J Cutan Med Surg*. 2013;17(6):47–54. <https://doi.org/10.2310/7750.2013.13072>
13. Leszczynski J, Lebski I. Anal warts (condylomata acuminata)-current issues and treatment modalities. *Adv Clin Exp Med*. 2014;23(2):307–311. <https://doi.org/10.17219/acem/37083>
14. Delany-Moretwe S, Kelley KF, James S, et al. Human papillomavirus vaccine introduction in South Africa: Implementation lessons from an evaluation of the national school-based vaccination campaign. *Global Health: Sci Pract*. 2018;6(3):425–438. <https://doi.org/10.9745/ghspp-d-18-00090>
15. Ngcobo NJ, Burnett RJ. Human papillomavirus vaccination acceptance and hesitancy in South Africa: Research and policy agenda. *S Afr Med J*. 2019;109(1):13–15. <https://doi.org/10.7196/samj.2018.v109i1.13723>
16. Mbulawa ZZ, Marais DJ. Impact of human immunodeficiency virus on the natural history of human papillomavirus genital infection in South African men and women. *J Infect Dis*. 2012;206(1):15–27. <https://doi.org/10.1093/infdis/jis299>
17. Clifford GM, Franceschi S, Keiser O, et al. Immunodeficiency and the risk of cervical intraepithelial neoplasia 2/3 and cervical cancer: A nested case-control study in the Swiss HIV cohort study. *Int J Cancer*. 2016;138(7):1732–1740. <https://doi.org/10.1002/ijc.29913>
18. Mbulawa ZZ, Johnson LF, Marais DJ, et al. Increased alpha-9 human papillomavirus species viral load in human immunodeficiency virus-positive women. *BMC Infect Dis*. 2014;14(1):1–7. <https://doi.org/10.1186/1471-2334-14-51>
19. McDonald AC, Tergas. Distribution of human papillomavirus genotypes among HIV-positive and HIV-negative women in Cape Town, South Africa. *Front Oncol*. 2014;4:48. <https://doi.org/10.3389/fonc.2014.00048>
20. Teixeira MF, Sabidó M, Leturiondo AL, et al. High risk human papillomavirus prevalence and genotype distribution among women infected with HIV in Manaus, Amazonas. *Viol J*. 2018;15(1):1–2. <https://doi.org/10.1186/s12985-018-0942-6>
21. Mbulawa ZZA, Somdyala NI. High human papillomavirus prevalence among females attending high school in the Eastern Cape Province of South Africa. *PLoS One*. 2021;16(6):e0253074. <https://doi.org/10.1371/journal.pone.0253074>
22. Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *PLoS Med*. 2007;6(7):e1000097. <https://doi.org/10.1093/ptj/89.9.873>
23. Bouvard V, Baan R, Straif K, et al. A review of human carcinogens--Part B: Biological agents. *The Lancet. Oncology*. 2009;10(4):321. [https://doi.org/10.1016/s1470-2045\(09\)70096-8](https://doi.org/10.1016/s1470-2045(09)70096-8)
24. Schiffman M, Clifford G. Classification of weakly carcinogenic human papillomavirus types: Addressing the limits of epidemiology at the borderline. *Infect Agents Canc*. 2009;4(1):1–8. <https://doi.org/10.1186/1750-9378-4-8>
25. Clifford GM, Gallus S, Herrero R, et al. Worldwide distribution of human papillomavirus types in cytologically normal women in the International Agency for Research on Cancer HPV prevalence surveys: A pooled analysis. *Lancet*. 2005;366(9490):991–998. [https://doi.org/10.1016/s0140-6736\(05\)67069-9](https://doi.org/10.1016/s0140-6736(05)67069-9)
26. Smith JS, Melendy A. Age-specific prevalence of infection with human papillomavirus in females: A global review. *J Adolesc Health*. 2008;43(4):S5–e10. <https://doi.org/10.1016/j.jadohealth.2008.07.009>
27. Ebrahim S, Mndende XK, Kharsany AB, et al. High burden of human papillomavirus (HPV) infection among young women in KwaZulu-Natal, South Africa. *PLoS One*. 2016;11(1):e0146603. <https://doi.org/10.1371/journal.pone.0146603>
28. Clifford G, Franceschi S. HPV in sub-Saharan Africa. *Papillomavirus Rep*. 2005;16(5):322–326. <https://doi.org/10.1179/095741905x4908>
29. Ari Ş, Arikian M. Next-generation sequencing: Advantages, disadvantages, and future. In: Khalid RH, Güzin T & Hüseyin T, editors. *Trends and applications*. Cham: Springer, 2016; p. 109–135.
30. Nilyanimit P, Chansaenroj J. Comparison of four human papillomavirus genotyping methods: Next-generation sequencing, INNO-LiPA, electrochemical DNA Chip, and nested-PCR. *Ann Lab Med*. 2018;38(2):139–146. <https://doi.org/10.3343/alm.2018.38.2.139>
31. Barzon L, Militello V, Lavezzo E, et al. Human papillomavirus genotyping by 454 next generation sequencing technology. *J Clin Virol*. 2011;52(2):93–97. <https://doi.org/10.1016/j.jcv.2011.07.006>
32. Flores-Miramontes MG, Torres-Reyes LA, Alvarado-Ruiz L, et al. Human papillomavirus genotyping by linear array and next-generation sequencing in cervical samples from Western Mexico. *Viol J*. 2015;12(1):1. <https://doi.org/10.1186/s12985-015-0391-4>
33. Meiring TL, Salimo AT, Coetzee B, et al. Next-generation sequencing of cervical DNA detects human papillomavirus types not detected by commercial kits. *Viol J*. 2012;9(1):164. <https://doi.org/10.1186/1743-422x-9-164>
34. Ardhaoui M, Ennaifer E, De Matos Salim AC, et al. Nested PCR followed by NGS: Validation and application for HPV genotyping of Tunisian cervical samples. *PLoS One*. 2021;16(8):e0255914. <https://doi.org/10.1371/journal.pone.0255914>
35. Mbulawa ZZ, Johnson LF. The impact of human immunodeficiency virus on human papillomavirus transmission in heterosexually active couples. *J Infect*. 2013;67(1):51–58. <https://doi.org/10.1016/j.jinf.2013.03.009>
36. Mbulawa ZZ, Coetzee D, Marais DJ, et al. Genital human papillomavirus prevalence and human papillomavirus concordance in heterosexual couples are positively associated with human immunodeficiency virus coinfection. *J Infect Dis*. 2009;199(10):1514–1524. <https://doi.org/10.1086/598220>
37. Clifford GM, Goncalves MA. Human papillomavirus types among women infected with HIV: A meta-analysis. *AIDS*. 2006;20(18):2337–2344. <https://doi.org/10.1097/01.aids.0000253361.63.578.14>
38. Akarolo-Anthony SN, Al-Mujtaba M, Famooto AO, et al. HIV associated high-risk HPV infection among Nigerian women. *BMC Infect Dis*. 2013;13(1):1–6. <https://doi.org/10.1186/1471-2334-13-521>
39. Kombe AJ, Li B, Zahid A, et al. Epidemiology and burden of human papillomavirus and related diseases, molecular pathogenesis, and vaccine evaluation. *Front Public Health*. 2020;8:552028. <https://doi.org/10.3389/fpubh.2020.552028>
40. De Sanjose S, Quint WG, Alemany L, et al. Human papillomavirus genotype attribution in invasive cervical cancer: A retrospective cross-sectional worldwide study. *Lancet Oncol*. 2010;11(11):1048–1056. [https://doi.org/10.1016/S1470-2045\(10\)70230-8](https://doi.org/10.1016/S1470-2045(10)70230-8)
41. Guan P, Howell-Jones R, Li N, et al. Human papillomavirus types in 115 789 HPV-positive women: A meta-analysis from cervical infection to cancer. *Int J Cancer*. 2012;131(10):2349–2359. <https://doi.org/10.1002/ijc.27485doi:10.1002/ijc.27485>
42. Tjalma WA, Fiander A, Reich O, et al. HERACLES/SCALE Study Group: Differences in human papillomavirus type distribution in high-grade cervical intraepithelial neoplasia and invasive cervical cancer in Europe. *Int J Cancer*. 2013;132(4):854–867. <https://doi.org/10.1002/ijc.2748>
43. Joura EA, Pils S. Vaccines against human papillomavirus infections: Protection against cancer, genital warts or both? *Clin Microbiol Infect*. 2016;22:S125–S127. <https://doi.org/10.1016/j.cmi.2016.12.017>
44. Di Resta C, Ferrari M. Next generation sequencing: From research area to clinical practice. *Electron J Int Fed Clin Chem Lab Med*. 2018;29(3):215–220.
45. Wolday D, Derese M, Gebressellasse S, et al. HPV genotype distribution among women with normal and abnormal cervical cytology presenting in a tertiary gynecology referral clinic in Ethiopia. *Infect Agents Canc*. 2018;13(1):1–8. <https://doi.org/10.1186/s13027-018-0201-x>
46. Muderris T, Afsar I. HPV genotype distribution among women with normal and abnormal cervical cytology in Turkey. *Revista Española de Quimioterapia*. 2019;32(6):516.
47. Obeid DA, Almatrouk SA. Human papillomavirus epidemiology in populations with normal or abnormal cervical cytology or cervical cancer in the Middle East and North Africa: A systematic review and meta-analysis. *J Infect Public Health*. 2020;13(9):1304–1313. <https://doi.org/10.1016/j.jiph.2020.06.012>
48. Strickler HD, Burk RD, Fazzari M, et al. Natural history and possible reactivation of human papillomavirus in human immunodeficiency virus-positive women. *J Natl Cancer Inst*. 2005;97(8):577–586. <https://doi.org/10.1093/jnci/dji073>
49. Coghill AE, Shiels MS. Elevated cancer-specific mortality among HIV-infected patients in the United States. *J Clin Oncol*. 2015;33(21):2376. <https://doi.org/10.1200/jco.2014.59.5967>
50. Dryden-Peterson S, Bvochora-Nsingo M, Suneja G, et al. HIV infection and survival among women with cervical cancer. *J Clin Oncol*. 2016;34(31):3749–3757. <https://doi.org/10.1200/jco.2016.67.9613>
51. Sankaranarayanan R, Budukh AM. Effective screening programmes for cervical cancer in low-and middle-income developing countries. *Bull World Health Organ*. 2001;79:954–962.
52. Choi YJ, Park JS. Clinical significance of human papillomavirus genotyping. *J Gynecol Oncol*. 2016;27(2):e21. <https://doi.org/10.3802/jgo.2016.27.e21>
53. Saslow D, Carolyn D, Runowicz CD, et al. American Cancer Society guideline for the early detection of cervical neoplasia and cancer. *CA Cancer J Clin*. 2002;52(6):342–362. <https://doi.org/10.3322/canjclin.52.6.342>
54. Jones J, Saleem A, Rai N, et al. Human papillomavirus genotype testing combined with cytology as a 'test of cure' post treatment: The importance of a persistent viral infection. *J Clin Virol*. 2011;52(2):88–92. <https://doi.org/10.1016/j.jcv.2011.06.021>
55. Paraskevaides E, Arbyn M, Sotiriadis A, et al. The role of HPV DNA testing in the follow-up period after treatment for CIN: A systematic review of the literature. *Cancer treatment reviews*. 2004;30(2):205–211. <https://doi.org/10.1016/j.ctrv.2003.07.008>
56. Snyman LC, Dreyer G. The Vaccine and Cervical Cancer Screen project 2 (VACCS 2): Linking cervical cancer screening to a two-dose HPV vaccination schedule in the South-West District of Tshwane, Gauteng, South Africa. *S Afr Med J*. 2015;105(3):191–194. <https://doi.org/10.7196/samj.8888>
57. Department of Health (Western Cape Government). Vaccinations. [cited 2022 Jan 05]. Available from: <https://www.westerncape.gov.za/general-publication/hpv-vaccinations>
58. Moodley I, Tathiah N. High uptake of Gardasil vaccine among 9-12-year old schoolgirls participating in an HPV vaccination demonstration project in KwaZulu-Natal, South Africa. *S Afr Med J*. 2013 103(5):318–321. <https://doi.org/10.7196/samj.6414>
59. Lefkowitz ES, Kelly KM. Correlates of human papillomavirus vaccination among female university students. *Women Health*. 2014;54(6):487–501. <https://doi.org/10.1080/03630242.2014.903552>
60. Danis K, Georgakopoulou T. Socioeconomic factors play a more important role in childhood vaccination coverage than parental perceptions: A cross-sectional study in Greece. *Vaccine*. 2010;28(7):1861–1869. <https://doi.org/10.1016/j.vaccine.2009.11.078>
61. Liu G, Sharma M. HIV-positive women have higher risk of human papilloma virus infection, precancerous lesions, and cervical cancer. *Aids*. 2018;32(6):795–808. <https://doi.org/10.1097/qad.0000000000001765>

62. Hidalgo-Tenorio C, Gil-Anguita C. ART is key to clearing oncogenic HPV genotypes (HR-HPV) in anal mucosa of HIV-positive MSM. *PLoS One*. 2019;14(10):e0224183. <https://doi.org/10.1371/journal.pone.0224183>
63. Taku O, Mbulawa ZZ. Distribution of Human Papillomavirus (HPV) genotypes in HIV-negative and HIV-positive women with cervical intraepithelial lesions in the Eastern Cape Province, South Africa. *Viruses*. 2021;13(2):280. <https://doi.org/10.3390/v13020280>
64. Saidu R, Kuhn L, Tergas A, et al. Performance of Xpert HPV on self-collected vaginal samples for cervical cancer screening among women in South Africa. *J Lower Genital Tract Dis*. 2021;25(1):15. <https://doi.org/10.1097/igt.0000000000000575>
65. Taku O, Brink A, Meiring TL, et al. Detection of sexually transmitted pathogens and co-infection with human papillomavirus in women residing in rural Eastern Cape, South Africa. *Peer J*. 2021;9:e10793. <https://doi.org/10.7717/peerj.10793>
66. Chamuso R, Ramesar R, Kaambo E, et al. Age, absolute CD4 count, and CD4 percentage in relation to HPV infection and the stage of cervical disease in HIV-1-positive women. *Medicine*. 2020;99(9):e19273. <https://doi.org/10.1097/md.00000000000019273>
67. Taku O, Meiring TL, Gustavsson I, et al. Acceptability of self-collection for human papillomavirus detection in the Eastern Cape, South Africa. *PLoS One*. 2020;15(11):e0241781. <https://doi.org/10.1371/journal.pone.0241781>
68. Kuhn L, Saidu R, Boa R, et al. Clinical evaluation of modifications to a human papillomavirus assay to optimise its utility for cervical cancer screening in low-resource settings: A diagnostic accuracy study. *Lancet Global Health*. 2020;8(2):e296–304. [https://doi.org/10.1016/s2214-109x\(19\)30527-3](https://doi.org/10.1016/s2214-109x(19)30527-3)
69. Johnson LG, Saidu R, Mbulawa Z, et al. Selecting human papillomavirus genotypes to optimize the performance of screening tests among South African women. *Canc Med*. 2020;9(18):6813–6824. <https://doi.org/10.1002/cam4.3329>
70. Taku O, Businge CB, Mdaka ML, et al. Human papillomavirus prevalence and risk factors among HIV-negative and HIV-positive women residing in rural Eastern Cape, South Africa. *Int J Infect Dis*. 2020;95:176–182. <https://doi.org/10.1016/j.ijid.2020.02.051>
71. Van De Wijgert JH, Gill AC, Chikandiwa A, et al. Human papillomavirus infection and cervical dysplasia in HIV-positive women: Potential role of the vaginal microbiota. *AIDS*. 2020;34(1):115–125. <https://doi.org/10.1097/qad.0000000000002381>
72. Chikandiwa A, Kelly H, Sawadogo B, et al. Prevalence, incidence and correlates of low risk HPV infection and anogenital warts in a cohort of women living with HIV in Burkina Faso and South Africa. *PLoS One*. 2018;13(5):e0196018. <https://doi.org/10.1371/journal.pone.0196018>
73. Mbulawa ZZ, Van Schalkwyk C, Hu NC, et al. High human papillomavirus (HPV) prevalence in South African adolescents and young women encourages expanded HPV vaccination campaigns. *PLoS One*. 2018;13(1):e0190166. <https://doi.org/10.1371/journal.pone.0190166>
74. Mbatha JN, Taylor M, Kleppa E, et al. High-risk human papillomavirus types in HIV-infected and HIV-uninfected young women in KwaZulu-Natal, South Africa: Implications for vaccination. *Infect Dis*. 2017;49(8):601–608. <https://doi.org/10.1080/23744235.2017.1312513>
75. Mbatha JN, Galappaththi-Arachchige HN, Mtshali A, et al. Self-sampling for human papillomavirus testing among rural young women of KwaZulu-Natal, South Africa. *BMC Res Notes*. 2017;10(1):1–8. <https://doi.org/10.1186/s13104-017-3045-3>
76. Kelly HA, Sawadogo B, Chikandiwa A, et al. Epidemiology of high-risk human papillomavirus and cervical lesions in African women living with HIV/AIDS: Effect of anti-retroviral therapy. *AIDS*. 2017;31(2):273–285. <https://doi.org/10.1097/qad.0000000000001301>
77. Kelly HA, Ngou J, Chikandiwa A, et al. Associations of Human Papillomavirus (HPV) genotypes with high-grade cervical neoplasia (CIN2+) in a cohort of women living with HIV in Burkina Faso and South Africa. *PLoS One*. 2017;12(3):e0174117. <https://doi.org/10.1371/journal.pone.0174117>
78. Dylla L, Abar B. Human papillomavirus clustering patterns among HIV-infected and HIV-uninfected adolescent females in South Africa. *J AIDS HIV Res*. 2017;9(10):202. <https://doi.org/10.5897/jahr2017.0445>
79. Rad A, Sørbye SW, Dreyer G, et al. HPV types in cervical cancer tissue in South Africa: A head-to-head comparison by mRNA and DNA tests. *Medicine*. 2017;96(47):e8752. <https://doi.org/10.1097/md.00000000000008752>
80. Adler DH, Wallace M, Bennie T, et al. Cumulative impact of HIV and multiple concurrent human papillomavirus infections on the risk of cervical dysplasia. *Adv Virol*. 2016;2016:7310894. <https://doi.org/10.1155/2016/7310894>
81. Segondy M, Kelly H, Magooa MP, et al. Performance of care HPV for detecting high-grade cervical intraepithelial neoplasia among women living with HIV-1 in Burkina Faso and South Africa: HARP study. *Br J Canc*. 2016;115(4):425–430. <https://doi.org/10.1038/bjc.2016>
82. Van Aardt MC, Dreyer G. Oncogenic and incidental HPV types associated with histologically confirmed cervical intraepithelial neoplasia in HIV-positive and HIV-negative South African women. *S Afr Med J*. 2016;106(6):617–622. <https://doi.org/10.7196/samj.2016.v106i6.10335>
83. Mbulawa ZZ, Wilkin TJ, Goeieman B, et al. Xpert human papillomavirus test is a promising cervical cancer screening test for HIV-seropositive women. *Papillomavirus Res*. 2016;2:56–60. <https://doi.org/10.1016/j.pvr.2016.02.004>
84. Pappasavvas E, Surrey LF, Glencross DK, et al. High-risk oncogenic HPV genotype infection associates with increased immune activation and T cell exhaustion in ART-suppressed HIV-1-infected women. *Oncoimmunology*. 2016;5(5):e1128612. <https://doi.org/10.1080/2162402x.2015.1128612>
85. Mbatani N, Adams T, Van Wijk L, et al. Performance of an human papillomavirus test in samples from women with histopathologically confirmed invasive cervical cancer. *J Lower Genital Tract Dis*. 2016;20(2):151–153. <https://doi.org/10.1097/igt.0000000000000183>
86. Mbulawa ZZ, Coetzee D. Human papillomavirus prevalence in South African women and men according to age and human immunodeficiency virus status. *BMC Infect Dis*. 2015;15(1):1. <https://doi.org/10.1186/s12879-015-1181-8>
87. Van Aardt MC, Dreyer G, Pienaar HF, et al. Unique human papillomavirus-type distribution in South African women with invasive cervical cancer and the effect of human immunodeficiency virus infection. *Int J Gynecol Canc*. 2015;25(5):919–925. <https://doi.org/10.1097/igc.0000000000000422>
88. Tayib S, Allan B. Human papillomavirus genotypes and clinical management of genital warts in women attending a colposcopy clinic in Cape Town, South Africa. *S Afr Med J*. 2015;105(8):679–684. <https://doi.org/10.7196/samjnew.7890>
89. Zeier MD, Botha MH, Engelbrecht S, et al. Combination antiretroviral therapy reduces the detection risk of cervical human papilloma virus infection in women living with HIV. *AIDS*. 2015;29(1):59–66. <https://doi.org/10.1097/qad.0000000000000512>
90. Adler D, Wallace M, Bennie T, et al. High risk human papillomavirus persistence among HIV-infected young women in South Africa. *Int J Infect Dis*. 2015;33:219–221. <https://doi.org/10.1016/j.ijid.2015.02.009>
91. Lebelo RL, Bogers JJ, Thys S, et al. Detection, genotyping and quantitation of multiple hpv infections in South African women with cervical squamous cell carcinoma. *J Med Virol*. 2015;87(9):1594–1600. <https://doi.org/10.1002/jmv.24132>
92. Ngou J, Gilham C, Omar T, et al. Comparison of analytical and clinical performances of the digene HC2 HPV DNA assay and the INNO-LiPA HPV genotyping assay for detecting high-risk HPV infection and cervical neoplasia among HIV-positive African women. *J Acquir Immune Defic Syndr*. 2015;68(2):162–168. <https://doi.org/10.1097/qai.0000000000000428>
93. Ameur A, Meiring TL, Buniks I, et al. Comprehensive profiling of the vaginal microbiome in HIV positive women using massive parallel semiconductor sequencing. *Scientific Reports*. 2014;4(1):1–7. <https://doi.org/10.1038/srep04398>
94. Adler DH, Wallace M, Bennie T, et al. Cervical dysplasia and high-risk human papillomavirus infections among HIV-infected and HIV-uninfected adolescent females in South Africa. *Infect Dis Obstet Gynecol*. 2014;2014:498048. <https://doi.org/10.1155/2014/498048>
95. Adler D, Laher F, Wallace M, et al. High rate of multiple concurrent Human Papillomavirus infections among HIV-uninfected South African Adolescents. *J Immunol Tech Infect Dis*. 2013;2(1):1000106.
96. Adler DH, Laher F, Lazarus E, et al. A viable and simple self-sampling method for human papillomavirus detection among South African adolescents. *J Immunol Tech Infect Dis*. 2013;2:3. <https://doi.org/10.4172/2329-9541.1000113>
97. Denny L, Adewole I, Anorlu R, et al. Human papillomavirus prevalence and type distribution in invasive cervical cancer in sub-Saharan Africa. *Int J Canc*. 2014;134(6):1389–1398. <https://doi.org/10.1002/ijc.28425>
98. Richter K, Becker P. Age-specific prevalence of cervical human papillomavirus infection and cytological abnormalities in women in Gauteng Province, South Africa. *S Afr Med J*. 2013;103(5):313–317. <https://doi.org/10.7196/samj.6514>
99. Ngou J, Magooa MP, Gilham C, et al. Comparison of care HPV and hybrid capture 2 assays for detection of high-risk human Papillomavirus DNA in cervical samples from HIV-1-infected African women. *J Clin Microbiol*. 2013;51(12):4240–4242. <https://doi.org/10.1128/jcm.02144-13>
100. Dols JA, Reid G, Kort R, et al. PCR-based identification of eight lactobacillus species and 18 hr-HPV genotypes in fixed cervical samples of South African women at risk of HIV and BV. *Diagn Cytopathol*. 2012;40(6):472–477. <https://doi.org/10.1002/dc.21786>
101. Dols JA, Reid G, Brown JM, et al. HPV type distribution and cervical cytology among HIV-positive Tanzanian and South African women. *Int Sch Res Not*. 2012;2012:514146. <https://doi.org/10.5402/2012/514146>
102. McDonald AC, Denny L. Distribution of high-risk human papillomavirus genotypes among HIV-negative women with and without cervical intraepithelial neoplasia in South Africa. *PLoS One*. 2012;7(9):e44332. <https://doi.org/10.1371/journal.pone.0044332>
103. De Vuyst HD, Ndirangu G, Moodley M, et al. Prevalence of human papillomavirus in women with invasive cervical carcinoma by HIV status in Kenya and South Africa. *Int J Canc*. 2012;131(4):949–955. <https://doi.org/10.1002/ijc.26470>
104. Taylor S, Wang C. A comparison of human papillomavirus testing of clinician-collected and self-collected samples during follow-up after screen and treat. *Int J Canc*. 2011;129(4):879–886. <https://doi.org/10.1002/ijc.25731>
105. Auvert B, Marais D. High-risk human papillomavirus is associated with HIV acquisition among South African female sex workers. *Infect Dis Obstet Gynecol*. 2011;2011:692012. <https://doi.org/10.1155/2011/692012>
106. Wang C, Wright TC. Rapid rise in detection of human papillomavirus (HPV) infection soon after incident HIV infection among South African women. *J Infect Dis*. 2011;203(4):479–486. <https://doi.org/10.1093/infdis/jiq083>
107. Mbulawa ZZ, Marais DJ. Influence of human immunodeficiency virus and CD4 count on the prevalence of human papillomavirus in heterosexual couples. *Journal of general virology*. 2010;91(12):3023–3031. <https://doi.org/10.1099/vir.0.020669-0>
108. Firnhaber C, Van Le H, Pettifor A, et al. Association between cervical dysplasia and human papillomavirus in HIV seropositive women from Johannesburg South Africa. *Canc Causes Contr*. 2010;21(3):433–443. <https://doi.org/10.1007/s10552-009-9475-z>
109. Firnhaber C, Zungu K, Levin S, et al. Diverse and high prevalence of human papillomavirus associated with a significant high rate of cervical dysplasia in human immunodeficiency virus-infected women in Johannesburg, South Africa. *Acta Cytol*. 2009;53(1):10–17. <https://doi.org/10.1159/000325079>

110. Moodley JR, Constant D, Hoffman M, et al. Human papillomavirus prevalence, viral load and pre-cancerous lesions of the cervix in women initiating highly active antiretroviral therapy in South Africa: A cross-sectional study. *BMC Canc*. 2009;9(1):1–8. <https://doi.org/10.1186/1471-2407-9-275>
111. Said HM, Ahmed K. HPV genotypes in women with squamous intraepithelial lesions and normal cervixes participating in a community-based microbicide study in Pretoria, South Africa. *J Clin Virol*. 2009;44(4):318–321. <https://doi.org/10.1016/j.jcv.2009.02.001>
112. Richter KL, Van Rensburg EJ. Human papilloma virus types in the oral and cervical mucosa of HIV-positive South African women prior to antiretroviral therapy. *J Oral Pathol Med*. 2008;37(9):555–559. <https://doi.org/10.1111/j.1600-0714.2008.00670.x>
113. Marais DJ, Passmore JA. Cervical and oral human papillomavirus types in HIV-1 positive and negative women with cervical disease in South Africa. *J Med Virol*. 2008;80(6):953–959. <https://doi.org/10.1002/jmv.21166>
114. Marais DJ, Constant D, Allan B, et al. Cervical human papillomavirus (HPV) infection and HPV type 16 antibodies in South African women. *J Clin Microbiol*. 2008;46(2):732–739. <https://doi.org/10.1128/jcm.01322-07>
115. Allan B, Marais DJ. Cervical human papillomavirus (HPV) infection in South African women: Implications for HPV screening and vaccine strategies. *J Clin Microbiol*. 2008;46(2):740–742. <https://doi.org/10.1128/jcm.01981-07>
116. Jones HE, Allan BR, Van De Wijgert JH, et al. Agreement between self-and clinician-collected specimen results for detection and typing of high-risk human papillomavirus in specimens from women in Gugulethu, South Africa. *J Clin Microbiol*. 2007;45(6):1679–1683. <https://doi.org/10.1128/jcm.02369-06>
117. Kay P, Soeters R. High prevalence of HPV 16 in South African women with cancer of the cervix and cervical intraepithelial neoplasia. *J Med Virol*. 2003;71(2):265–273. <https://doi.org/10.1002/jmv.10479>
118. Kuhn L, Denny L. Human papillomavirus DNA testing for cervical cancer screening in low-resource settings. *J Natl Canc Inst*. 2000;92(10):818–825. <https://doi.org/10.1093/jnci/92.10.818>
119. Wright Jr TC, Denny L. HPV DNA testing of self-collected vaginal samples compared with cytologic screening to detect cervical cancer. *JAMA*. 2000;283(1):81–86. <https://doi.org/10.1001/jama.283.1.81>
120. Ramesar JE, Dehaeck CM. Human papillomavirus in normal cervical smears from Cape Town. *S Afr Med J*. 1996;86(11).
121. Williamson AL, Brink NS. Typing of human papillomaviruses in cervical carcinoma biopsies from Cape Town. *J Med Virol*. 1994;43(3):231–237. <https://doi.org/10.1002/jmv.1890430307>
122. Cooper K, Herrington CS. Integration of human papillomavirus types 16 and 18 in cervical adenocarcinoma. *J Clin Pathol*. 1992;45(5):382–384. <https://doi.org/10.1136/jcp.45.5.382>
123. Cooper K, Herrington CS. In situ human papillomavirus (HPV) genotyping of cervical intraepithelial neoplasia in South African and British patients: Evidence for putative HPV integration in vivo. *J Clin Pathol*. 1991;44(5):400–405. <https://doi.org/10.1136/jcp.44.5.400>
124. Cooper K, Herrington CS. In situ evidence for HPV 16, 18, 33 integration in cervical squamous cell cancer in Britain and South Africa. *J Clin Pathol*. 1991;44(5):406–409. <https://doi.org/10.1136/jcp.44.5.406>
125. Williamson AL, Dehaeck CM. Typing of human papillomaviruses in cervical intraepithelial neoplasia grade 3 biopsies from Cape Town. *J Med Virol*. 1989;28(3):146–149. <https://doi.org/10.1002/jmv.1890280307>